



**Summertime Acute Heat Illness in U.S. Emergency
Departments from 2006 through 2010:
Analysis of a Nationally Representative Sample**

Jeremy J. Hess, Shubhayu Saha, and George Luber

<http://dx.doi.org/10.1289/ehp.1306796>

Received: 13 March 2013

Accepted: 16 June 2014

Advance Publication: 17 June 2014

Summertime Acute Heat Illness in U.S. Emergency Departments from 2006 through 2010: Analysis of a Nationally Representative Sample

Jeremy J. Hess,^{1,2,3} Shubhayu Saha,¹ and George Luber¹

¹Climate and Health Program, Division of Environmental Hazards and Health Effects, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA; ²Department of Emergency Medicine, Emory University School of Medicine, Atlanta, Georgia, USA; ³Department of Environmental Health, Rollins School of Public Health at Emory University, Atlanta, Georgia, USA

Address correspondence to Jeremy Hess, Emory University Department of Emergency Medicine, 49 Jesse Hill Jr Drive, Atlanta, GA 30303 USA. Telephone: 404-251-8851. E-mail: jhess@emory.edu

Running title: Heat Illness in US Emergency Departments

Acknowledgments: The authors would like to thank Anne Elixhauser, Dana Flanders, Debra Houry, Josephine Malilay, Rebecca Noe, and Stephen Pitts for helpful comments.

Competing financial interests: Dr. Hess has served as a scientific consultant to the Natural Resources Defense Council and to Lumenetec, LLC. Neither of the other authors has any actual or potential competing financial interests to declare.

Abstract

Background: Patients with acute heat illness present primarily to emergency departments (EDs), yet little is known regarding these visits.

Objective: To describe acute heat illness visits to US EDs from 2006-2010 and identify factors associated with hospital admission or death-in-the-ED.

Methods: We extracted ED case-level data from the Nationwide Emergency Department Sample (NEDS) for 2006-10, defining cases as ED visits from May-September with any heat illness diagnosis (ICD-9-CM 992.0-992.9). We correlated visit rates and temperature anomalies analyzed demographics and ED disposition, identified risk factors for adverse outcomes, and examined ED case fatality rates (CFR).

Results: There were 326,497 (95% CI: 308,372-344,658) cases, with 287,875 (88.2%) treated-and-released, 38,392 (11.8%) admitted, and 230 (0.07%) died-in-the-ED. Heat illness diagnoses were first-listed in 68%. 74.7% had heat exhaustion, 5.4% heat stroke. Visit rates were highly correlated with annual temperature anomalies (correlation coefficient 0.90, $p=0.037$). Treat-and-release rates were highest for younger adults (26.2/100,000/year), while hospitalization and death-in-the-ED rates were highest for older adults (6.7 and 0.03/100,000/year respectively); all rates were highest in rural areas. Heat stroke had an ED CFR of 99.4/10,000 (78.7-120.1) visits and was diagnosed in 77.0% of deaths. Adjusted odds of hospital admission or death-in-the-ED were higher among elders, males, urban and low income residents, and those with chronic conditions.

Conclusions: Heat illness presented to the ED frequently, with highest rates in rural areas. Case definitions should include all diagnoses. Visit rates were correlated with temperature anomalies.

Heat stroke had a high ED CFR. Males, elders, and the chronically ill were at greatest risk of admission or death-in-the-ED. Chronic disease burden exponentially increased this risk.

Introduction

Background

Heat has various effects on the human body, ranging from several well-recognized acute heat illness syndromes (Kilbourne 1997), exacerbations of chronic lung and cardiovascular disease (Kovats and Hajat 2008), and multiple other syndromes associated with dehydration (Josseran et al. 2009). Acute heat illnesses are a subset of hyperthermia syndromes (Stitt 1979) resulting primarily from diminished heat dissipation capacity, classically from environmental exposure, though exertion may be a factor. Acute heat illness severity ranges from mild, e.g. heat rash, cramps, and fatigue, to moderate, e.g. heat syncope and heat exhaustion, to life-threatening, e.g. heat stroke. Heat stroke is defined as a core temperature above 40°C and evidence of neurological dysfunction (Knochel and Reed 1994). Mild to moderate heat illness is reversible with prompt treatment. Heat stroke, however, has a high case fatality rate, 51% at 28 days to 71% at two years in one cohort, and survivors have significant sequelae (Argaud et al. 2007). While there is insufficient evidence surrounding the best resuscitative measures (Smith 2005), outcomes are clearly improved by early intervention (Bouchama et al. 2007).

While heat-related mortality is extensively studied, not as much is known about heat-related morbidity generally or acute heat illness specifically (Ye et al. 2012). In the US, acute heat illness presents most commonly to the emergency department (ED), and over 80% of acute heat illness hospitalizations come through the ED, twice the proportion for other illnesses (Merrill et al. 2008). ED volume increases during periods of intense heat: A study of the 2006 California heat wave, for instance, found that the rate ratio for acute heat illness (defined as International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM) (Centers for Disease Control and Prevention 2013) codes of 992) was 6.3 for ED visits and 10.2 for hospitalizations

compared with the reference period (Knowlton et al. 2009). To date, however, studies of heat illness in the ED have been limited in scope, focusing on particular events, demographic, or subsets of ED visits.

There are several compelling reasons for examining acute heat illness in the ED. First, it is likely a good indicator of trends for other, more prevalent diseases affected by heat, as evidenced in the 2006 California heat wave (Knowlton et al. 2009). Second, many vulnerable groups use the ED disproportionately (Pitts et al. 2008), including elders, the poor, the chronically ill, and those with mental health concerns (Ishigami et al. 2008). Third, the ED is an appropriate setting for surveillance (Josseran et al. 2009) and tertiary prevention. Fourth, emergency physicians frequently clarify and advocate for appropriate preventive measures (Kellermann and Todd 1996). Fifth, ED heat illness visits may become more prevalent as the frequency of record heat increases (Meehl et al. 2009). In general, climate change has implications for heat illness as part of routine occupational and recreational activities (Maloney and Forbes 2011) and is likely to impact emergency providers disproportionately (Hess et al. 2009).

Despite the importance of the ED in managing acute heat illness, there are no comprehensive national estimates of ED visits for heat that include both fatal and nonfatal cases and are based on all ED diagnoses for a patient's visit. Moreover, the available research does not clarify whether factors associated with presentation to the ED for heat illness are the same as those for heat-related deaths. Different risk profiles for morbidity and mortality based on disease pathogenesis (e.g. heat-related cerebrovascular injuries are more likely to be rapidly fatal than heat-related respiratory disease exacerbations) have been observed elsewhere (Kovats et al. 2004) and there may be implications for prevention programming and surveillance.

Objectives

We had three objectives with this analysis. The first was to estimate the burden of summertime acute heat illness using a nationally-representative sample of ED patients to generate population-based rates for acute heat illness ED visits. The second was to characterize acute heat illness case profiles across time and different ED dispositions. Our third objective was to identify demographic factors and comorbid conditions associated with adverse outcomes of hospital admission or death in the ED among patients with acute heat illness.

Methods

Study design

This study was an analysis of five recent years of data (2006-2010) from the Nationwide Emergency Department Sample (NEDS) of the Healthcare Cost and Utilization Project (HCUP) (<http://www.ahrq.gov/research/data/hcup/index.html>), an annual stratified sample of hospital-based EDs sponsored by the Agency for Healthcare Research and Quality (AHRQ) (Steiner et al. 2002). The NEDS nationally representative database is constructed using a 20% stratified sample of US hospital-based EDs from states contributing data to HCUP. Detailed information regarding sampling methods is available online at the AHRQ website (AHRQ 2011) and elsewhere (Owens et al. 2010). The NEDS has been used for multiple studies of various conditions seen commonly in EDs and is among the few nationally representative ED datasets available (Owens et al. 2010).

Assessment of possible exposure bias in NEDS sample

Our first step was to check for potential exposure bias related to the NEDS sampling scheme, as the NEDS database was constructed with data from select states (there were 24 partner states in 2006; five were added through 2009, and 28 participated in 2010; see Supplemental Material,

Table S1 for additional details). We checked if anomalously high temperature in the NEDS states were similar to non-NEDS states over the study period, on the supposition that the prevalence of ED visits for acute heat illness in selected states could not be nationally representative if there were systematic differences in exposure. From the National Climatic Data Center (NCDC) website (NCDC 2013; <http://www.ncdc.noaa.gov/cag/>), we obtained historic temperature information for the lower 48 states. For each study year we compared the difference between average daily maximum temperature for May 1 to September 30 to the 30-year normal (1971-2000) for each state. We then compared the average differences in maximum temperature for the NEDS states with the non-NEDS states and tested whether these differences were statistically significant (see Supplemental Material, Table S2). The tests show that the temperature patterns were statistically different between the two groups only for the year 2009, with the anomaly in non-NEDS states greater than NEDS states.

Data collection and management

Our next step was to complete the NEDS data use agreements and download ED visit data from AHRQ. Data for national population estimates were also downloaded from the AHRQ website (Agency for Healthcare Research and Quality (AHRQ) 2012). Insurance data were downloaded from the US Census Bureau website (U.S. Census Bureau 2013). Data included demographics, insurance status (primary billed insurance), urban-rural designation of case's county of residence [adapted from the NCHS Urban-Rural Classification Scheme for Counties (http://www.cdc.gov/nchs/data_access/urban_rural.htm)], ED diagnoses (first-listed and other diagnoses available in the record of ED visit), and disposition (treat-and-release or discharge, admission to the same hospital, transfer to another facility, and died-in-the-ED). We analyzed the data using SAS version 9.3 (SAS Institute, Inc., Cary, NC).

Our outcomes of interest were any ED diagnosis of acute heat illness as designated by the ICD-9-CM codes 992.0 to 992.9 and/or with an injury code (e-code) of E900 (accident or injury from exposure to excessive environmental heat); hospital admission; and death-in-the-ED. To increase the likelihood that cases were representative of environmental heat exposure, we defined cases as patients presenting to the ED during summertime months (May 1 to September 30 inclusive). We excluded cases with an e-code indicating heat exposure in a man-made environment (E900.1).

We used the Clinical Classifications Software (CCS) created by HCUP for ICD-9-CM to collapse the large array of diagnoses into a smaller number of clinically meaningful categories (<http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>) and used these categories in our regression analyses.

HCUP also provides a binary Chronic Condition Indicator (CCI), which we used to classify ICD-9-CM diagnosis codes as chronic or not. We then categorized the number of chronic conditions and created an index of chronic disease burden (categories included one, two, three, and four or more). We used this index in a series of multiple regressions to evaluate the contribution of comorbid chronic disease on adjusted odds of hospital admission or death in the ED among all ED visits in the sample, among all cases except those with heat stroke, and among heat stroke cases.

As a secondary data analysis, the study was exempted from institutional review.

Data analysis

To pursue our first and second objectives we estimated total counts and population-based rates of ED visits for acute heat illness by ED disposition. We then disaggregated these counts based on (i) whether or not heat illness was listed as the first diagnosis for the visit; (ii) by year of visit and

demographic and geographic factors; and (iii) different heat illness diagnoses. We used the statistical methodology prescribed by AHRQ (AHRQ 2012) to incorporate the complex survey design and utilize appropriate weights to generate national counts of ED episodes. We also used prescribed AHRQ methodology to estimate national rates for specific variables (AHRQ, 2012). As ambient temperature anomalies varied across time (see Supplemental Material Table S2), we used Spearman's rho to evaluate the association between average annual temperature anomaly in NEDS states and annual ED heat illness visit rate.

To pursue our third objective we conducted a series of regressions on the entire sample and on subsets defined by specific heat illness diagnoses. Since there was a gradient in severity of outcomes, we first considered ordinal logistic regression, but we decided against that approach as model diagnostics indicated that the proportional odds assumption was violated. We also estimated a polytomous regression model with the three ED dispositions, but were circumspect of these results given the relatively small number of died-in-the-ED cases. We elected to present these results in Supplemental Material Table S3. Our final approach was to perform logistic regression on a binary outcome variable created by combining the hospitalization and death-in-the-ED outcomes into one group. For the logistic regression we compared this composite outcome with the treat-and-release disposition. Independent variables included case-level demographic information, health insurance status, median income for case's residential ZIP code, urban classification of county of residence, and CCS category of comorbid disease as described above.

Our last set of regressions used chronic disease burden category as described above as a predictor variable and examined adjusted odds of the composite outcome in the entire sample and in cases with heat stroke and any acute heat illness except heat stroke, controlling for the

same variables described in the analyses above. Our hypothesis in this analysis was that adjusted odds of the composite outcome would be positively correlated with chronic disease burden.

To evaluate the ED case fatality rate (CFR) for heat stroke (the rate at which cases with heat stroke died in the ED), we compared with ED CFR for heat stroke cases with the ED CFR for the cases with any heat illness diagnosis as well as the ED CFR for all ED visits from 2006-2010 inclusive, stratified by CCS.

Results

Estimates of summertime acute heat illness ED visits

Table 1 presents estimates of mean case counts and annualized population-based rates stratified by first-listed and secondary diagnosis and ED disposition. There were an estimated 326,497 (95% CI 308,372-344,658) cases in the study period (94% of all heat illness visits in the sample occurred between May 1 and September 30). Dividing by the number of years in the study period yields an average of 65,299 ED visits for summertime acute heat illness per year, an average rate of 21.5 visits/100,000 population/year. According to estimates provided by HCUP (<http://hcupnet.ahrq.gov/HCUPnet.jsp>), there were 625 million ED visits in the United States from 2006-2010. Using this as the denominator and assuming no repeat visits for heat illness, this suggests an approximate average of 5/10,000 ED visits were for summertime heat illness during the study period.

About 88% of cases were treated and released, 12% were admitted to the hospital, and a small number died in the ED (Table 1). 68% of the cases had a first-listed diagnosis of heat illness and among them, 91% were treated and released. In the 32% of cases with a secondary diagnosis of heat illness, 81% were treated and released. The difference in percentage of treat-and-release

disposition between cases with primary and secondary heat illness diagnoses was statistically significant ($p < 0.001$). However, the demographic, socioeconomic, insurance, and comorbid condition profiles of the two groups were not statistically different (results available upon request).

Acute heat illness case profiles across time and ED dispositions

Table 2 shows univariate associations between demographic and other factors and ED disposition for all cases of Acute Heat Illness related ED visits from May 1 to September 30 during the study period. Both counts as well as rates per 100,000 population per year are presented. Missing data were rare; the most commonly missing data were for income (3% of the observations), rurality (1%), and type of health insurance (11%).

Annual counts and rates showed significant interannual variation, with highest counts and rates (apart from the rate of death-in-the-ED) in 2010. Annual temperature anomalies for NEDS states are shown in Supplemental Material Table S2. As shown in Figure 1, there is a significant correlation between annual temperature anomalies and annual population-based rate for ED heat illness visits.

In regards to demographic variables, the 18-45 age group had the highest treat-and-release rate, while the 65+ age group had the highest rates of hospital admission and death-in-the-ED. Males were more likely than females to have heat illness across all dispositions. The highest rates for treat-and-release and hospital admissions were in the South; the highest rate of death-in-the-ED was in the West. Across all ED dispositions, the highest rates were found among people in the bottom income quartile. Areas designated as rural and micropolitan had higher rates across all ED dispositions compared to the areas designated as ‘metropolitan.’ With respect to health

insurance, the uninsured population had the highest rate among the treat-and-release visits, while cases with Medicare had the highest rates of hospital admission and death-in-the-ED.

Factors associated with adverse outcomes of hospital admission or death-in-the-ED

Counts of heat illness diagnoses stratified by ED disposition are presented in Table 3. 74.7% of cases were diagnosed with heat exhaustion (ICD9 codes 992.3 – 992.5), 5.4% with heat stroke (ICD9 code 992.0) and 3.7% with non-specific heat illness (ICD9 code 992.9). Among those with heat stroke, 62.0% of cases were either admitted (28.3% of all admissions) or died in the ED (77.0% of all death-in-the-ED cases). The risk of fatality was much lower among heat exhaustion cases, 90.6% of which were treated and released when considered altogether.

Table 4 presents results from the logistic regression examining factors associated with the composite adverse outcome (hospital admission or death-in-the-ED) among all cases. Adjusted odds ratios of hospitalization or death-in-the-ED relative to treat-and-release are presented as the model controls for a wide range of factors – demographic, geographic, rural-urban indicator, health insurance type, chronic conditions and diagnosis of heat illness. The >65 group, males, residents of large metropolitan areas, the uninsured, residents of ZIP codes with low median household incomes, and cases with heat stroke had higher odds of hospitalization or death-in-the-ED than treatment-and-release. There was relatively little meaningful significant regional or interannual variation compared with the significant rate differences in Table 2. Adjusted odds of hospital admission or death-in-the-ED were increased by all chronic conditions in some cases substantially: chronic hematologic conditions increased adjusted odds nine-fold.

While the small number of died-in-the-ED cases make us circumspect, we present the results of the polytomous logistic regression in Supplemental Material, Table S3. In this analysis, adjusted

odds of death-in-the-ED were highest for cases aged 0-17, males, Midwestern residents, residents of low-income ZIP codes, those with Medicaid, and those with chronic circulatory conditions.

Since the adjusted odds ratio of hospitalization or death-in-the-ED among heat stroke cases was 23.86 (22.35-25.46) (Table 4), we separately analyzed if these risk factors were different among this subgroup (Supplemental Material, Table S4 and Table S5). Among heat stroke cases, younger patients had higher adjusted odds of the composite outcome than non-heat stroke cases; adjusted odds were also higher for heat stroke cases in large metropolitan areas, though the observed urban-rural gradient was the same. In both groups, cases from the Midwest had the lowest adjusted odds, though the difference was more marked for heat stroke cases. Other trends were similar between the two groups.

Table 5 shows the results of the ED CFR analysis. The CFR for all acute heat illness diagnoses is comparable to those of the other CCS groups. The CFR for heat stroke, 99.4/10,000 (95% CI 78.7-120.1) ED visits, is significantly higher than the other groups, the highest of which is 50.7/10,000 for circulatory disease.

Figure 2 shows the results of the logistic regression analyses examining the association between the index of chronic disease burden and adjusted odds of the composite outcome (hospital admission or death) for three categories of ED patients: those with any heat illness except heat stroke, those with heat stroke, and all ED visits. In each analysis we found a sharp increase in the relationship between chronic conditions and adjusted odds of the composite outcome. The curve slopes varied, and were lowest for heat stroke cases and highest for all ED visits. Put differently,

higher chronic disease burden was more strongly associated with odds of hospitalization or death-in-the-ED for all ED visits than for heat stroke cases.

Discussion

Our findings contribute to the burgeoning literature on heat and morbidity generally and, in particular, ED visits for acute heat illness. Our analysis suggests that, for the years studied, the NEDS is likely to provide reasonable national estimates of ED visits for summertime acute heat illness. Our results also suggest that acute heat illness is a substantial concern, and that heat exhaustion is the most common presentation. Our findings provide new evidence regarding the relatively high rates of acute heat illness in rural populations, the high ED CFR of heat stroke, and the dramatic role of age and chronic conditions on the likelihood of hospital admission or death-in-the-ED. Our results also suggest that odds of death-in-the-ED are highest among the young, poor, Midwestern males, though this requires further study. Finally, our results also appear to capture inter-annual variability in dose-response, with higher visit rates in anomalously warm years (see Figure 1 and Supplemental Material Table S2).

It is important to note that our estimates are for acute heat illness only and not for associated conditions whose prevalence may be affected by heat exposure (i.e. the broader category of heat-associated illness). While incidence of acute heat illness can rise dramatically during extreme heat events, incidence of several heat-associated illnesses also rises, and acute heat illness is a relatively small proportion of the total increased disease burden. In the 2006 California heat wave, for instance, acute heat illness cases made up only 13.2% of the excess ED visits (Knowlton et al. 2009). There was increased incidence in other disease categories, particularly electrolyte imbalance, nephritis, and renal failure, exclusive of an associated acute heat illness

diagnosis. Thus, if patterns observed during heat waves can be applied to other summertime periods, our estimates likely represent only a fraction of the total burden of heat-related ED visits. Without exposure data, however, we cannot assess the validity of this claim.

The lack of exposure and other data also complicate interpretation of some of our other findings. The regional differences in died-in-the-ED rates among patients with heat illness are one example, and may relate to regional differences in exposure, illness severity, or admission practices, or to variability resulting from the relatively small sample size. The relatively high rate of heat illness in rural areas, which may reflect differences in occupational exposures in certain settings or to differential prevalence in air conditioning availability and usage, is another example.

There is only one prior national estimate of the incidence of ED visits for acute heat illness against which we can compare our findings (Sanchez et al. 2010). Our estimates of heat exhaustion incidence, male predominance, and the age distribution of heat illness cases are all relatively consistent with theirs. The main difference is the overall incidence of heat illness among ED patients. They estimated that there were 20,775 heat illness ED visits a year from 2001-2004, a third of our estimate of for years 2006-2010. Part of the difference can be attributed to differences in case definition, as cases in that study were nonfatal visits with documented environmental heat exposure and first-listed diagnosis of heat injury. In our sample, limiting the case definition to nonfatal cases with first-listed diagnosis of heat illness (Table 1) yielded 221,640 cases for the study period or an average of 44,328 such cases annually, roughly double their estimate. Other possible causes for this difference are unclear. There may be bias in either or both of the estimates, or differential exposure among the sampled populations. While temperatures have been gradually warming during this time, there has not been a significant

difference in the frequency of heat waves between the two study periods (Environmental Protection Agency 2010). Regional temperature differences would presumably have a larger impact on estimates using the National Electronic Injury Surveillance System-All Injury Program (NEISS-AIP, http://www.healthindicators.gov/Resources/DataSources/NEISS-AIP_88/Profile), the data Sanchez et al. used, which collects data from 66 hospitals, than the NEDS, which samples roughly 1,000, but as investigators are blinded to hospital identities it is difficult to assess this explanation. Finally, the other study's strict inclusion criteria surely play some role and likely resulted in an underestimate, as the authors note.

We can also compare a subset of our findings with another national estimate of rates of ED visits for hyperthermia among Medicare beneficiaries in 2004-5 (Noe et al. 2012). Here, too, our estimate is higher. They estimate a rate of 12.0 ED hyperthermia visits/100,000 population/year and 5.3 hospital admissions (not all from the ED) for heat illness/100,000 population/year with an admission proportion (the proportion of visits that result in hospital admission) of 11%. For cases with Medicare we estimate a rate of 24.6 ED visits/100,000 population/year, an admission rate from the ED of 6.9/100,000 population/year, and an admission proportion of 28.0% (Table 2). The differences are most likely due to inclusion criteria: Noe et al. included only cases with a first-listed diagnosis of hyperthermia, whereas we included cases if acute heat illness was one of any listed. Heat illness was listed first in 67.9% of our cases (Table 1) and this could account for the bulk of the difference, highlighting the importance of including secondary diagnoses in epidemiological studies of ED visits for heat illness. The remainder of the difference may result from inter-annual variability in visit rates and differences in the databases used for the respective analyses.

As might be expected, our findings suggest that heat illness is more prevalent among young adults and that hospitalization is more prevalent among older adults. Perhaps surprisingly, death-in-the-ED is equally prevalent in the young and in elders. Differences in exposure and/or susceptibility may be responsible for these patterns. Younger people are presumably more exposed through occupational and recreational activities and have greater physiologic resilience, suggesting that a larger proportion of their heat illness presentations are less severe. This could, paradoxically, be the explanation for the relatively high prevalence of death-in-the-ED among cases aged 0-17, in the event that younger people are able to compensate in the earlier stages of heat illness and present later in the illness course when treatment is less effective. That elders are more likely to be admitted to the hospital is potentially a function of higher exposure, to the extent that elders may have less capacity to recognize and reverse heat illness at earlier stages, or it may be related to their greater burden of underlying chronic disease. The higher prevalence of death-in-the-ED among elders is presumably driven by physical frailty and underlying comorbid disease. Our data (Table 4) do demonstrate an association between chronic disease burden and complications of heat illness, suggesting that physiologic reserve is at least partially responsible, but this explanation is not exclusive.

That increasing chronic disease burden is associated with increased adjusted odds of hospital admission or death in the ED is unsurprising. What is perhaps more surprising is that the relative increase in risk conferred by additional chronic diseases was lower for heat stroke cases than for all heat illness cases and for any ED visits in general. This finding is somewhat difficult to interpret. It may be because heat stroke is a serious condition that frequently warrants admission in-and-of itself, and that comorbid conditions carry less weight in the admission decision calculus in such cases. But the fact that other diagnoses are more commonly listed first,

suggesting greater concern about these other conditions, belies this interpretation. Alternatively, exposures may be different for those with heat stroke compared with other acute heat illnesses. Evaluation of the true relationship will likely require exposure data or perhaps other methods such as a chart review.

That some chronic diseases appear to confer greater risk is not surprising given the pathophysiology and social determinants of heat illness. The finding that chronic mental health disorders increase risk is well established, for instance (Page et al. 2012). The strongly increased risk associated with chronic hematologic disease, however, is a new finding, though an association between heat illness and sickle cell anemia (anemia is the most common chronic hematologic disease reported in the NEDS) is established (Smith et al. 2003). While the association between chronic circulatory disease and death-in-the-ED is established, its strength in our polytomous regression is remarkable and, like the other findings, requires confirmation and further study. Overall, our findings support the established public health tenet of importance of targeting prevention messages at patients with chronic conditions.

Our findings reinforce previous work regarding public health programming related to extreme heat. The large burden of heat illness in a working-age population may relate to occupational exposure though this warrants further study; regardless, prevention should target at-risk workers (Bernard and McGeehin 2004). The higher odds of adverse outcomes among elders suggest that efforts should target these populations as well (without creating a false sense of security in other groups), again consistent with prior findings (Luber and McGeehin 2008). We can also infer from our results that EDs should expect that those with the highest risk of dying in the ED may be the young, while increases in hospital admission are most likely to be for older patients with multiple chronic conditions.

These findings provide additional information for those interested in relating shifts in the frequency and severity of heat hazards, population susceptibility factors, and outcomes such as death in the ED and hospital admission. Our results may also have utility in projections of climate change impacts on heat morbidity or impacts on emergency services as well as more comprehensive estimates of costs associated with extreme heat events. To project such impacts, however, exposure-outcome associations will be needed, not just to characterize acute heat illness burdens but also to characterize increases in other heat-associated illness, as discussed above. These associations cannot be derived without linking exposure and outcome data.

Our findings speak to the strengths of the NEDS as a dataset, including a large sample size, national representation, and availability of up to 15 ED diagnoses per visit. These strengths allow for a high level of statistical power in generating national estimates for small subpopulations and patients with multiple diagnoses, including chronic diseases. Comparison with other national ED data sources found that NEDS is among the best for detailed examination of diagnoses (Owens et al. 2010). Compared with the National Hospital Ambulatory Medical Care Survey (NHAMCS, <http://www.cdc.gov/nchs/ahcd.htm>), the other leading national ED dataset, NEDS captures a higher number of diagnoses and a larger proportion of elderly patients (Owens et al. 2010). Compared with other databases for injury-related ED visits, which our outcomes include, the NEDS and the NEISS-AIP provide comparable estimates of injuries (though both provide lower injury estimates than NHAMCS) (Owens et al. 2010).

Despite these strengths, however, our study is subject to several limitations, most importantly the inability to link outcomes with exposure data. In the NEDS, ED visit date and location are restricted to protect patients' and hospitals' confidentiality. We used visit month as a proxy and excluded exposures of known man-made etiology. While this approach captured 94% of heat

illness cases, some could be unrelated to summertime heat exposure. Similarly, it is reasonable to note that the external validity of these findings depends on the study years 2006 to 2010 being representative not only of trends in non-NEDS states, which our analyses confirm, but also of recent temperature trends. Several of the years we studied were anomalously warm compared with climate normals, but this is in keeping with recent temperature trends (Hansen et al. 2012). Some other potential limitations arising from the use of the NEDS data are also worth mentioning. The data included in the NEDS are based on the coding by providers and coders, and surveillance and research were not the intended use of these data. Moreover, admission diagnoses are also included for the subset of patients admitted to the hospital in the NEDS sample. These aspects of the survey's design may introduce several biases into our results that bear mention.

Two potential biases may have resulted in overestimates related to severe heat illness. First, as there is no information on the inter-rater reliability of ED diagnoses of heat illness, it is possible that providers who treat patients with heat exhaustion are more likely to diagnose heat stroke than another acute heat illness if there are associated complications, even if there is no change in mental status. This could lead to an overestimate of heat stroke, though it is difficult to estimate the magnitude of its potential impact. Second, diagnoses are coded and arranged to maximize reimbursement for the care provided, not to facilitate epidemiologic research. This could introduce selection bias wherein severe illness is diagnosed more frequently. If present, this would also result in an overestimate of severe heat illness. We feel this is unlikely given that heat stroke was more often a secondary diagnosis, suggesting that the billing provider more often felt that heat stroke was secondary to another, more significant primary process.

Two other potential biases are worth noting. First, there may have been detection bias from differential recognition of heat illness depending on perceptions of the weather, e.g. during the latter part of a declared heat wave. Conversely, there may have been underrecognition of heat illness during shorter periods of more routine weather. If present, such bias could result in over or underestimation of heat illness, and so it is difficult to assess its likely direction or magnitude. Second, providers probably did not code for all relevant chronic conditions. This would have resulted in an underestimate of the role of individual chronic disease categories in heat illness, particularly in treat-and-release visits, as those who were admitted presumably had a more exhaustive enumeration of comorbid and chronic conditions. To the extent that patients tend to have multiple chronic conditions, this underreporting may conversely have resulted in a diffuse overestimate of the observed associations with chronic disease burden. It is difficult to estimate the magnitude of these potential biases.

With these caveats in mind, we conclude that the NEDS provides a more comprehensive estimate of heat illness presentations to US EDs than other nationally representative ED datasets and provides useful insights into the role of chronic conditions in the clinical course of heat illness patients.

Conclusion

Analysis of this nationally representative sample of ED visits for summertime acute heat illness demonstrates that its incidence was higher than anticipated, particularly in rural areas, low income communities, and among the uninsured. Most ED visits for heat illness were for less severe forms. Our analysis reveals patterns of heat illness similar to prior estimates but suggests a larger burden of disease. Population-based rates were highly correlated with average temperature anomalies in our five-year sample, suggesting ED visits for heat illness may be a

reasonable surveillance indicator, though evaluation with a longer time series or more granular exposure information is warranted. Our findings provide new insight into dynamics surrounding heat stroke, which was more frequently diagnosed among older patients and those with multiple chronic conditions. Heat stroke cases had a high ED case fatality rate, and risk of death-in-the-ED appears highest in young, poor, males, and Midwesterners. Chronic illness significantly increased risk of hospital admission or death-in-the-ED, and the number of chronic conditions was associated with increased risk. The relative increase in risk is greater for all heat diagnoses than heat stroke, highlighting the severity of this condition. Our findings reinforce previous work on heat preparedness, and may be useful in projecting heat morbidity associated with climate change. Future research could explore exposure-outcome associations by linking ED visits with temperature exposures and explore linkages between ED diagnoses and management with clinical outcomes and costs.

References

- Agency for Healthcare Research and Quality (AHRQ). 2012. Population Denominator Data for Use with the HCUP Databases (Updated with 2009 Population Data), Report # 2010-02. Available: http://hcup-us.ahrq.gov/reports/methods/2010_02.pdf [accessed 7/25/2012].
- AHRQ. 2011. National Emergency Department Sample (NEDS) of the Healthcare Cost and Utilization Project (HCUP). Available: www.hcup-us.ahrq.gov/nedsoverview.jsp [accessed 7/25/2012].
- Applegate W, Runyan J, Brasfield L, Williams M, Konigsberg C, Fouche C. 1981. Analysis of the 1980 heat wave in Memphis. *J Am Geriatr Soc* 29:337-342.
- Argaud L, Ferry T, Le Q-H, Marfisi A, Ciorba D, Achache P, et al. 2007. Short- and Long-term Outcomes of Heatstroke Following the 2003 Heat Wave in Lyon, France. *Arch Intern Med* 167(20):2177-2183.
- Bernard SM, McGeehin MA. 2004. Municipal heat wave response plans. *Am J Public Health* 94(9):1520-1522.
- Bouchama A, Dehbi M, Chaves-Carballo E. 2007. Cooling and hemodynamic management in heatstroke: practical recommendations. *Crit Care* 11(3):R54.
- Centers for Disease Control and Prevention. 2013. International Classification of Diseases, 9th Revision, Clinical Modification. Available: <http://www.cdc.gov/nchs/icd/icd9cm.htm> [accessed 12 November 2013].
- Environmental Protection Agency. 2010. Climate Change Indicators in the United States. Washington, D.C.:Environmental Protection Agency. Available: http://epa.gov/climatechange/indicators/jpdfs/ClimateIndicators_full.pdf [accessed 7/25/2012].
- Hansen J, Sato M, Ruedy R. 2012. Perception of climate change. *Proceedings of the National Academy of Sciences* 109(37):E2415-E2423.
- Hess J, Heilpern K, Davis T, Frumkin H. 2009. Climate Change and Emergency Medicine: Impacts and Opportunities. *Academic Emergency Medicine* 16:782-794.
- Ishigami A, Hajat S, Kovats RS, Bisanti L, Rognoni M, Russo A, et al. An ecological time-series study of heat-related mortality in three European cities. 2008. *Environmental Health*. 7(5):1-7, doi:10.1186/1476-069X-7-5.

- Josseran L, Caillere N, Brun-Ney D, Rottner J, Filleul L, Brucker G, et al. 2009. Syndromic surveillance and heat wave morbidity: a pilot study based on emergency departments in France. *BMC Medical Informatics & Decision Making* 9(14):1-9, doi:10.1186/1472-6947-9-14.
- Kellermann AL, Todd KH. 1996. Killing heat. *New England Journal of Medicine* 335(2):126-127.
- Kilbourne E. 1997. Heat waves and hot environments. In: *The Public Health Consequences of Disasters*, (Noji E, ed). New York:Oxford University Press, 245-269.
- Knochel J, Reed G. 1994. Disorders of heat regulation. In: *Maxwell & Kleeman's clinical disorders of fluid and electrolyte metabolism*, 5th ed, (Narrins R, ed). New York:McGraw-Hill, 1549-1590.
- Knowlton K, Rotkin-Ellman M, King G, Margolis H, Smith D, Solomon G, et al. 2009. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 117(1):61-67.
- Kovats RS, Hajat S. 2008. Heat stress and public health: a critical review. *Annual Review of Public Health* 29:41-55.
- Kovats RS, Hajat S, Wilkinson P. 2004. Contrasting patterns of mortality and hospital admissions during hot weather and heat waves in Greater London, UK. *Occupational & Environmental Medicine* 61(11):893-898.
- Luber G, McGeehin M. 2008. Climate Change and Extreme Heat Events. *Am J Prev Med* 35:429-436.
- Maloney S, Forbes C. 2011. What effect will a few degrees of climate change have on human heat balance? Implications for human activity. *International Journal of Biometeorology* 55(2):147-160.
- Meehl GA, Tebaldi C, Walton G, Easterling D, McDaniel L. 2009. Relative increase of record high maximum temperatures compared to record low minimum temperatures in the U.S. *Geophys Res Lett* 36(23):L23701.
- Merrill C, Miller M, Steiner C. 2008. Hospital Stays Resulting from Excessive Heat and Cold Exposure Due to Weather Conditions in U.S. Community Hospitals, 2005. Rockville, MD:Agency for Healthcare Research and Quality. Available: <http://www.ncbi.nlm.nih.gov/books/NBK56045/> [accessed 7/25/2012].

- National Climatic Data Center (NCDC). 2013. Climate at a Glance. Available: <http://www.ncdc.noaa.gov/cag/> [accessed 12 November 2013].
- Noe RS, Jin JO, Wolkin AF. 2012. Exposure to Natural Cold and Heat: Hypothermia and Hyperthermia Medicare Claims, United States, 2004–2005. *American Journal of Public Health* 102(4):e11-e18.
- Owens PL, Barrett ML, Gibson TB, Andrews RM, Weinick RM, Mutter RL. 2010. Emergency Department Care in the United States: A Profile of National Data Sources. *Annals of Emergency Medicine* 56(2):150-165.
- Page L, Hajat S, Kovats R, Howard L. 2012. Temperature-related deaths in people with psychosis, dementia and substance misuse. *Br J Psychiatry* 200:485-490.
- Pitts SR, Niska RW, Xu J, Burt CW, Pitts SR, Niska RW, et al. 2008. National Hospital Ambulatory Medical Care Survey: 2006 emergency department summary. *National health statistics reports*(7):1-38.
- Sanchez C, Thomas K, Malilay J, Annett J. 2010. Nonfatal Natural and Environmental Injuries Treated in Emergency Departments, United States, 2001–2004. *Fam Community Health* 33(1):3-10.
- Smith J. 2005. Cooling methods used in the treatment of exertional heat illness. *British Journal of Sports Medicine* 39:503-507.
- Smith WR, Coyne P, Smith VS, Mercier B. 2003. Temperature changes, temperature extremes, and their relationship to emergency department visits and hospitalizations for sickle cell crisis. *Pain Management Nursing* 4(3):106-111.
- Steiner C, Elixhauser A, Schnaier J. 2002. The healthcare cost and utilization project: an overview. *Effective clinical practice : ECP* 5(3):143-151.
- Stitt JT. 1979. Fever versus hyperthermia. *Federation proceedings* 38(1):39-43.
- U.S. Census Bureau. 2013. Health Insurance Historical Tables - HIB Series. Available: http://www.census.gov/hhes/www/hlthins/data/historical/HIB_tables.html [accessed 11/12/13 2013].
- Ye X, Wolff R, Yu W, Vaneckova P, Pan X, Tong S. 2012. Ambient Temperature and Morbidity: A Review of Epidemiological Evidence. *Environ Health Perspect* 120(1):19-28.

Table 1. Counts and rates^a (per 100,000 population per year) of acute heat illness (ICD-9 code = 992.xx) with 95% confidence intervals (CIs) by first-listed and secondary diagnosis and Emergency Department (ED) disposition^b.

ED disposition	First-listed diagnosis	Secondary diagnoses	Combined
Treat-and-release			
N	202829 (192015, 213643)	85046 (79493, 90599)	287875 (272134, 3036152)
rate	13.3	5.6	18.9
Admit-to-hospital			
N	18811 (17494, 20127)	19581 (18426, 20736)	38392 (36051, 40732)
rate	1.2	1.3	2.5
Died-in-the-ED			
N	99 (68, 129)	131 (86, 177)	230 (187, 274)
rate	0.007	0.009	0.015
Total			
N	221739 (209577, 233899)	104758 (98055, 111512)	326497 (308372, 344658)
rate	14.6	6.9	21.5

^aSum of total US population for 2006 to 2010 used as denominator; population estimates used for each year were: 309,349,689 (2010), 306,771,529 (2009), 304,093,966 (2008), 301,231,207 (2007), 298,379,912 (2006) (Source: http://hcup-us.ahrq.gov/reports/methods/2010_02.pdf). ^bEmergency Department visits between May 1 to September 30 in a calendar year.

Table 2. Descriptive statistics [n (rate)]^{a,b} of patients with acute heat illness (ICD9 = 992.0-992.9) by emergency department (ED) disposition (N = 326,497).^c

Variable	Treat and Release	Admit to Hospital	Death in ED
Year			
2006	61182 (20.5)	8214 (2.75)	99 (0.033)
2007	59641 (19.8)	7539 (2.5)	38 (0.013)
2008	48668 (16)	6416 (2.11)	-
2009	42517 (13.9)	5729 (1.87)	-
2010	75867 (24.5)	10494 (3.39)	57 (0.018)
Age groups			
0-17 years	44342 (12)	1430 (0.39)	102 (0.028)
18 - 45 years	147241 (26.2)	11432 (2.03)	-
46 - 65 years	67363 (17.1)	12634 (3.22)	69 (0.018)
> 65 years	28918 (14.9)	12896 (6.65)	59 (0.031)
Gender			
Male	197043 (25.5)	28591 (3.7)	186 (0.024)
Female	90676 (12.1)	9801 (1.31)	45 (0.006)
Hospital region			
Northeast	32130 (11.7)	3997 (1.46)	-
Midwest	64701 (19.5)	7365 (2.21)	-
South	146630 (26.2)	19406 (3.46)	80 (0.014)
West	44414 (12.6)	7623 (2.16)	117 (0.033)
Median zipcode income quartile			
Bottom quartile	91390 (23.9)	13372 (3.5)	103 (0.027)
Second quartile	81860 (21.6)	9957 (2.63)	55 (0.014)
Third quartile	61171 (16.2)	7613 (2.02)	30 (0.008)
Top quartile	45174 (11.9)	5656 (1.49)	-
Urban-rural classification			
Large metropolitan	112067 (13.8)	17546 (2.16)	114 (0.014)
Medium/small metropolitan	99258 (21.9)	12011 (2.65)	60 (0.013)
Micropolitan	43781 (29)	4948 (3.27)	30 (0.02)
Rural	29862 (29.7)	3340 (3.32)	-
Payment source^d			
Medicare	38427 (17.7)	14950 (6.88)	75 (0.035)
Medicaid	35713 (16.8)	3581 (1.68)	70 (0.033)
Private insurance	115112 (11.5)	10058 (1)	-
Uninsured	67062 (28.8)	6058 (2.6)	48 (0.021)

^aTotal number of observations under each category maybe different due to missing data. ^bRates are per 100,000 population per year; denominators were obtained from (AHRQ 2012). ^cData pooled for 2006-10; ED visits between 1 May - 30 September included; sample weights used.

^dHealth insurance coverage status 2006-10 obtained from (US Census Bureau 2013).

Table 3. Counts of acute heat illness diagnoses (ICD9 = 992.0-992.9) by emergency department (ED) disposition with row and column percents.^{a,b}

ICD-9-CM Heat Illness Diagnosis	Treat and Release (N)	Row ^c %	Column ^d %	Admit to Hospital (N)	Row %	Column %	Death in ED (N)	Row %	Column %
9920 - Heat Stroke	6889	38	2	11059	61	28	178	1	77
9921 - Heat Syncope	12812	86	4	2089	14	5			
9922 - Heat Cramps	17592	98	6	373	2	1			
9923 - Heat exhaustion (anhydrotic)	10841	79	4	2899	21	7			
9924 - Heat exhaustion (salt depletion)	1252	67	0	623	33	2			
9925 - Heat exhaustion (unspecified)	213659	91	73	19894	9	51	11	0	5
9926 - Heat Fatigue	2600	95	1	130	5	0	5	0	2
9927 - Heat Edema	629	99	0	9	1	0			
9928 - Other specified heat effects	16135	91	5	1585	9	4	11	0	5
9929 - Unspecified	11848	96	4	410	3	1	26	0	11

^aData pooled for 2006-10; ED visits between 1 May 30 September included; sample weights used. ^bPatients could have more than one diagnosis of acute heat illness during a visit. Total diagnoses count therefore is greater than the number of patients. ^cFor each specific heat diagnoses, Row % refers to the percentage of patients with different ED disposition. ^dFor each ED disposition, Column % refers to the percentage of patients with different heat diagnoses.

Table 4. Logistic regression - comparing odds ratios of hospital admission/death among patients with heat stroke (ICD9 = 992.0) and other acute heat illness diagnoses (ICD9 = 992.1 – 992.9) (N = 282743^{*}).^{a,b}

Variable	OR (95% CI)
Age groups	
0-17 yrs	0.4 (0.37,0.43)
18-45 yrs	0.45 (0.42,0.47)
45-65 yrs	0.59 (0.56,0.62)
65+ yrs (ref)	1
Urban-rural classification	
Large Metropolitan	1.36 (1.29,1.44)
Medium-Small Metropolitan	0.96 (0.91,1.02)
Micropolitan	0.93 (0.87,0.99)
Rural (ref)	1
Health insurance	
Medicare	1.01 (0.96,1.07)
Medicaid	0.88 (0.84,0.93)
Private	0.94 (0.9,0.98)
Uninsured (ref)	1
Median income for ZIP code	
Top quartile	0.76 (0.72,0.8)
Quartile 3	0.75 (0.72,0.79)
Quartile 2	0.83 (0.8,0.86)
Lowest quartile (ref)	1
Geographic regions	
Midwest	0.86 (0.82,0.91)
South	0.92 (0.88,0.97)
West	1 (0.94,1.06)
Northeast (ref)	1
Gender	
Male	1.64 (1.59,1.7)
Female (ref)	1
Associated chronic conditions	
Blood	9.05 (8.45,9.69)
Circulatory	2.44 (2.36,2.53)
Digestive	2.14 (2.02,2.26)
Endocrine	2.85 (2.76,2.95)
Genito-urinary	3.48 (3.25,3.72)
Mental	2.75 (2.66,2.84)
Muscular	2.43 (2.28,2.58)
Nervous	3.22 (3.05,3.39)
Respiratory	1.92 (1.83,2.02)
Cancer	2.66 (2.3,3.07)
No comorbid conditions (ref)	1
Heat diagnoses	
Heat stroke (992.0)	23.86 (22.35,25.46)
Heat syncope (992.1)	2.81 (2.6,3.04)
Heat exhaustion (992.3 - 992.5)	2.03 (1.93,2.14)
Other heat DX (ref)	1

Variable	OR (95% CI)
Year	
2010	0.73 (0.7,0.76)
2009	0.68 (0.65,0.72)
2008	0.79 (0.76,0.83)
2007	0.87 (0.83,0.91)
2006	1

^aData pooled for 2006-10; ED visits between 1 May 30 September included; sample weights used. ^bThe number of cases is smaller compared to Table 3 as some explanatory variables had missing variables.

Table 5. Comparing fatalities per 10,000 ED visits for acute heat illness and other illnesses based on HCUP Comorbidity Classification System (pooled data from 2006-10 from the entire NEDS sample).

HCUP Clinical classification	Rate per 10,000 ED Visits (95% CI)
Infectious and parasitic disease	7 (6.5,7.4)
Neoplasms	29.3 (27.9,30.6)
Endocrine, nutritional, metabolic diseases and immunity disorders	15.9 (15.1,16.7)
Diseases of blood and blood-forming organs	9.1 (8.6,9.6)
Mental illness	7.1 (6.7,7.5)
Diseases of nervous system and sense organs	5.8 (5.5,6.2)
Diseases of circulatory system	50.7 (48.5,52.9)
Diseases of respiratory system	13 (12.3,13.6)
Diseases of digestive system	5.8 (5.5,6.1)
Diseases of genitourinary system	8.7 (8.2,9.2)
Complications of pregnancy, childbirth and puerperium	0.7 (0.5,0.9)
Diseases of skin and subcutaneous tissue	1.9 (1.8,2.1)
Diseases of musculoskeletal and connective tissue	3.5 (3.2,3.7)
Congenital anomalies	12 (11.3,12.6)
Certain conditions originating in perinatal period	18.6 (17.4,19.7)
All acute heat illness diagnoses	8.9 (7.8,10)
Heat stroke	99.4 (78.7,120.1)

Figure legends

Figure 1. Annual population-based rates of ED visits for heat illness by average annual temperature anomalies for NEDS states. A trendline with 95% CI is included for reference. The Spearman's rho correlation coefficient is 0.90, $p=0.037$.

Figure 2. Adjusted odds ratios of hospital admission or death-in-the-ED for different degrees of chronic disease burden, stratified by all ED visits, all acute heat illness, and only heat stroke visit (ICD9 code 992.0). The index was calculated based on combining the Clinical Classification Software (CCS) and Chronic Condition Indicator (CCI) information provided in HCUP.

Figure 1




Figure 2

